

Sabrina Strickland, M.D.
Hospital for Special Surgery
535 East 70th Street
New York, NY 10021
(212) 606-1725

**Arthroscopic Capsular Release
Physical Therapy Protocol and Prescription**

Date:
Name:

Immediate Postoperative Period

- Sling for comfort only and as little as possible
- PROM without limits
- Emphasis on terminal range
- AAROM (wand, self-stretch)
- Ice 3-4x daily

7 Days-2 Weeks (Goal: Full PROM)

- Discontinue sling
- AAROM: add pulley and UBE for motion
- Isometrics for all shoulder motions within pain-free ROM
- AROM: impingement exercises without resistance (standing ABD, FE, supraspinatus exercises <90°, sidelying IR< ER; prone horizontal abduction, extension, flexion)
- Ice following exercises

3 Weeks (Goal: full AROM)

- PRE: hand weights or theraband resistance within pain-free ROM
- Impingement exercises
- Scapulothoracic exercises
 - Wall push-ups, supine punch-ups
 - Prone scapular retraction with horizontal abduction with IR and ER
 - Rowing; shrugs, press-ups
- Ice following exercises

4 Weeks (exercises should be pain-free)

- Progress PREs
- Add CKC exercises (step ups, BAPS, treadmill; STAB exercises)
- Add manual resistive exercises (PNF, rhythmic STAB, eccentrics)
- Add isokinetic rotation at neutral ABD
- Ice following exercises

6 Weeks (Full pain free ROM)

- Add plyometrics (medicine ball, theraband, plyoback)
- Resume sport specific activities (progression toward full activity)